

Deformations with section: Cotangent cohomology, flatness conditions and modular subgerms

Tobias Hirsch*

email: hirsch@math.tu-cottbus.de

BTU Cottbus, Institut für Mathematik,
Postfach 10 13 44, D-03046 Cottbus

Bernd Martin*

email: martin@math.tu-cottbus.de

BTU Cottbus, Institut für Mathematik,
Postfach 10 13 44, D-03046 Cottbus

February 1, 2008

Abstract

We study modular subspaces corresponding to two deformation functors associated to an isolated singularity X_0 : the functor Def_{X_0} of deformations of X_0 and the functor $Def_{X_0}^s$ of deformations with section of X_0 . After recalling some standard facts on the cotangent cohomology of analytic algebras and the general theory of deformations with section, we give several criteria for modularity in terms of the relative cotangent cohomology modules of a deformation. In particular it is shown that the modular strata for the functors Def_{X_0} and $Def_{X_0}^s$ of quasihomogeneous complete intersection singularities coincide. Flatness conditions for the first cotangent cohomology modules of the deformation functors under consideration are then compared.

Contents

Introduction	2
1 A Summary of cotangent cohomology	2
2 Deformations with section	7
3 Modular deformations	12
4 Flatness and modular deformations	20
References	25

*Partially supported by G. I. F.

Introduction

When trying to construct moduli spaces for analytic objects, one major difficulty results from the fact that, in general, semi-universal deformations of the objects under consideration are not universal. One approach to construct local moduli is the study of some kind of maximal universal locus in the base space of a semi-universal deformation. The corresponding notion of a modular deformation has been introduced by V. P. Palamodov in [Pal78] for complex spaces, later on in a formal context by O. A. Laudal in [Lau79].

In these notes we study the strictly local situation of germs of analytic spaces, i. e. deformations of singularities. This has been done before, see e. g. [Mar02a] and [Mar]. One obtains the following characterization of modular subspaces in the base S of a semi-universal deformation $\xi : X \rightarrow S$ of a singularity X_0 : $M \subseteq S$ is modular if and only if all vector fields of the special fibre can be lifted to vertical vector fields of the family ξ – a criterion that already appeared in [Pal78] in the context of compact complex spaces. For deformations of complete intersections and space curves, modularity of a subspace $M \subseteq S$ can also be interpreted as flattening stratum of the relative Tjurina-module $T^1(X/S)$ of the family. This can be used to actually compute non-trivial examples of modular strata using a new algorithm to determine local flattenings. The algorithm and its implementation in the computer algebra system SINGULAR, cf. [GPS01], are also explained in [Mar02a].

This article investigates the similar situation of deformations with section of X_0 , i. e. deformations $\xi : X \rightarrow S$ of X_0 together with a morphism $\sigma : S \rightarrow X$ such that $\xi \circ \sigma = id_S$. The first section is a brief review of the cotangent cohomology of analytic algebras and their morphisms which we will use extensively later on.

Section 2 collects the basic facts on deformations with section. In particular we state the results in Buchweitz' thesis [Buc81] on the cotangent cohomology of such deformations and on the construction of a semi-universal deformation with section, it is given explicitly for the case of complete intersections.

In section 3 we derive a variant of the Kodaira-Spencer sequence of the deformation ξ that also contains information on the given section. With this tool at hand we formulate and prove a criterion of modularity as liftability of vector fields on X_0 with values in the maximal ideal of \mathcal{O}_{X_0} to vertical vector fields of the family. As an application we prove that, for quasi-homogeneous complete intersections, the modular strata with respect to both deformation functors coincide, this being mainly a consequence of A. G. Alexandrov's description of the module of derivations $\text{Der}_{\mathbb{C}}(\mathcal{O}_{X_0})$ for this class of singularities in [Ale85].

In the closing section we give an interpretation of this criterion as flatness of the first cotangent cohomology module of X over S with coefficients in the kernel of $\sigma^* : \mathcal{O}_X \rightarrow \mathcal{O}_S$. We then compare flatness conditions for the zeroth and first relative cotangent cohomology modules. We finish by investigating the relationship between modular subspaces for both deformation functors and prove that both spaces coincide for deformations with singular section of hypersurfaces.

1 A Summary of cotangent cohomology

In this introductory section we summarize the facts about the cotangent cohomology of a morphism $(X, 0) \rightarrow (S, 0)$ of complex space germs or, equivalently, of morphisms $\mathcal{O}_{S,0} \rightarrow \mathcal{O}_{X,0}$ of analytic algebras that we will need later on. Since we are working exclusively in the category

(*Gan*) of germs of complex spaces (resp. the category (*Analg*) of analytic algebras), we usually omit the distinguished point and simply write X for a germ $(X, 0)$ of a complex space.

1.1 Cotangent cohomology of analytic algebras

We start by recalling the construction of the cotangent complex of a morphism $A \rightarrow B$ of analytic algebras and the definition and properties of the so-called cotangent cohomology functors $T^i(B/A, -)$. This is the complex-analytic analogue of Andre-Quillen cohomology (cf. [Qui70], [And74]). Details and proofs of the statements below can be found in [Pal76], [Fle78], [Buc81], [Pal82], for example.

Definition. Let $A \rightarrow B$ be a homomorphism of analytic algebras.

- (1) A *resolvent* R for B over A is a DG-algebra (a differential graded anticommutative analytic algebra) that is a free A -algebra, together with a surjective homomorphism $p : R \twoheadrightarrow B$ over A which is a quasiisomorphism of complexes, i. e. $H^n(R) = 0$ for $n < 0$ and $H^0(R) \simeq B$, and this isomorphism is induced by p .
- (2) The complex

$$\mathbb{L}_{B/A}^\bullet := \Omega_{R/A} \otimes_R B$$

is called the *cotangent complex* of B over A . Here $\Omega_{R/A}$ denotes the module of *Kähler differentials* of R over A .

- (3) For a B -module M we define the *cotangent cohomology modules* of B over A with values in M as

$$T^i(B/A, M) := \text{Ext}_B^i(\mathbb{L}_{B/A}^\bullet, M), \quad i \geq 0.$$

Remark. For the existence of a resolvent of B over A and the well-definedness of the functors T^i we refer to the papers cited above. In order to simplify notations we adopt to the conventions

$$T^i(B/A) := T^i(B/A, B), \quad T^i(B, M) := T^i(B/\mathbb{C}, M).$$

Additionally, if $X \rightarrow S$ is a morphism of space germs and M is an \mathcal{O}_X -module we also write

$$T^i(X/S, M) := T^i(\mathcal{O}_X/\mathcal{O}_S, M).$$

Proposition 1.1. The functors T^i have the following properties:

- (1) $T^0(B/A, M) \simeq \text{Der}_A(B, M)$.
- (2) If $A \rightarrow B$ turns B into a regular A -algebra, then $T^i(B/A, M) = 0$ for $i > 0$.
- (3) If $B \simeq A\{x\}/(f_1, \dots, f_k)$, where f_1, \dots, f_k form a regular sequence in $A\{x\}$, then $T^i(B/A, M) = 0$ for $i > 1$.
- (4) If $A \twoheadrightarrow B$ is a surjection with kernel I then $T^1(B/A, M) \simeq \text{Hom}_B(I/I^2, M)$.
- (5) If $T^1(B/A, \mathbb{C}) = 0$, then B is a regular A -algebra.
- (6) If $T^2(B/A, \mathbb{C}) = 0$, then $B \simeq A\{x\}/(f_1, \dots, f_k)$ with a regular sequence $f_1, \dots, f_k \in A\{x\}$.

The functorial properties of cotangent cohomology yield the following exact sequence:

Proposition 1.2. Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be a short exact sequence of B -modules. Then there is a long exact sequence in cohomology

$$0 \rightarrow T^0(B/A, M') \rightarrow T^0(B/A, M) \rightarrow T^0(B/A, M'') \rightarrow T^1(B/A, M') \rightarrow \dots$$

Furthermore, suppose we are additionally given a morphism $B \rightarrow C$, then there is an exact sequence

$$0 \rightarrow \mathbb{L}_{B/A}^\bullet \otimes_B C \rightarrow \mathbb{L}_{C/A}^\bullet \rightarrow \mathbb{L}_{C/B}^\bullet \rightarrow 0,$$

from which one derives:

Proposition 1.3. If $A \rightarrow B \rightarrow C$ are homomorphisms of analytic algebras and M is a C -module, then there is a long exact sequence in cohomology

$$0 \rightarrow T^0(C/B, M) \rightarrow T^0(C/A, M) \rightarrow T^0(B/A, M) \rightarrow T^1(C/B, M) \rightarrow \dots$$

In particular, combining Proposition 1.1 and Proposition 1.3 we get:

Proposition 1.4. Let $B \simeq A\{x\}/I$ be a quotient of a regular analytic A -algebra $A\{x\}$, and let M be a B -module. Then there is an exact sequence

$$0 \rightarrow \mathrm{Der}_A(B, M) \rightarrow \mathrm{Der}_A(A\{x\}, M) \rightarrow \mathrm{Hom}_B(I/I^2, M) \rightarrow T^1(B/A, M) \rightarrow 0.$$

Proof. Applying Proposition 1.3 to the homomorphisms $A \rightarrow A\{x\} \rightarrow B$ we get the exact sequence

$$0 \rightarrow T^0(B/A, M) \rightarrow T^0(A\{x\}/A, M) \rightarrow T^1(B/A\{x\}, M) \rightarrow T^1(B/A, M) \rightarrow 0,$$

using property (2) in Proposition 1.1 and the fact that $T^0(B/A\{x\}, M) \simeq \mathrm{Der}_{A\{x\}}(B, M) = 0$. Now the statement follows from the identities (1) and (4) in Proposition 1.1. \blacksquare

Remark. In fact, the sequence above results from applying $\mathrm{Hom}_B(-, M)$ to the so-called *conormal sequence*

$$I/I^2 \rightarrow \Omega_{A\{x\}/A} \otimes_{A\{x\}} B \rightarrow \Omega_{B/A} \rightarrow 0,$$

cf. [Eis95], for instance.

Finally, the following proposition describes the behaviour of the T^i -functors under base change:

Proposition 1.5. Let be given homomorphisms $A \rightarrow B$ and $A \rightarrow A'$ of analytic algebras, and set $B' := B \otimes_A A'$, i. e. we have a cocartesian diagram

$$\begin{array}{ccc} B & \longrightarrow & B' = B \otimes_A A' \\ \uparrow & & \uparrow \\ A & \longrightarrow & A'. \end{array}$$

If $A \rightarrow A'$ or $A \rightarrow B$ is flat, then there is a natural isomorphism

$$T^i(B/A, M) \simeq T^i(B'/A', M)$$

for any B' -module M and $i \geq 0$.

Remark. If, in the situation above, both $A \rightarrow A'$ and $B \rightarrow B'$ are flat, then there is also the identity $T^i(B/A, M) \otimes_A A' \simeq T^i(B'/A', M \otimes_A A')$ for any B -module M . If only $A \rightarrow A'$ is flat, we still have such a map, but in general this map needs not be an isomorphism.

So far we have introduced the notions and basic properties of the relative cotangent cohomology of a morphism $A \rightarrow B$ of analytic algebras; taking $A := \mathbb{C}$ we get the absolute cotangent cohomology of B . Now let X_0 be a germ of a complex space, then one has (cf. [Pal76, § 5], [Buc81, 2.4.4]) the following basic interpretation of $T^i(X_0)$ in terms of *deformations* of X_0 , i. e. flat maps $\xi : X \rightarrow S$ with $\xi^{-1}(0) \simeq X_0$ (here and subsequently we denote by D the one-point space $\{0\}$ with $\mathcal{O}_D \simeq \mathbb{C}[\varepsilon]/(\varepsilon^2)$):

- $T^0(X_0)$ is isomorphic to the module of infinitesimal automorphisms of X_0 , i. e. automorphisms of $X_0 \times D$ over D ,
- $T^1(X_0)$ coincides with the space of first-order infinitesimal deformations, i. e. the tangent space of the *deformation functor*

$$Def_{X_0}(S) := \{\text{isomorphism classes of deformations of } X_0 \text{ over } S\},$$

- $T^2(X_0)$ contains the obstructions of lifting deformations to higher order.

Analogous statements hold for the modules $T^i(X/S)$ that control relative deformations of X over S , given by a diagram

$$\begin{array}{ccc} X & \xrightarrow{i} & \mathcal{X} \\ \downarrow & & \downarrow \\ S & \xrightarrow{j} & S \times B \end{array} \quad \begin{array}{c} \searrow \xi \\ \nearrow pr_2 \end{array} \quad B.$$

1.2 The cohomology functors $T^i(f, -)$

In what follows we will also consider deformations of morphisms $f : X \rightarrow S$ of germs, their corresponding cotangent cohomology is given by functors $T^i(f, -)$ that we are now going to describe. Again, for details and the application of the following constructions (and their global counterparts) to deformations of morphisms of complex spaces, we refer to [Fle78].

Definition. Let A, B and C be analytic algebras such that B and C are A -algebras, and let $f : B \rightarrow C$ be an A -algebra homomorphism (for the applications we have in mind, we are of course mainly interested in the case $A = \mathbb{C}$). Choose resolvents R (resp. S) of B (resp. C) over A that are compatible with f , i. e. there is a commutative diagram

$$\begin{array}{ccc} R & \longrightarrow & S \\ \downarrow & & \downarrow \\ B & \xrightarrow{f} & C \end{array}$$

of DG-algebras over A .

(1) Define

$$\mathbb{L}_f^\bullet := (\mathbb{L}_{B/A}^\bullet, \mathbb{L}_{C/A}^\bullet).$$

Formally, this is a complex in the following category \mathcal{C} : \mathcal{C} is the (abelian) category whose objects are tripels (M_1, M_2, h) of a B -module M_1 , a C -module M_2 and a homomorphism $h : M_1 \rightarrow M_2$ over $f : B \rightarrow C$. $\text{Mor}_{\mathcal{C}}((M_1, M_2, h), (M'_1, M'_2, h'))$ consists of all pairs (g_1, g_2) of a B -module homomorphism $g_1 : M_1 \rightarrow M'_1$ and a C -module homomorphism $g_2 : M_2 \rightarrow M'_2$ such that $g_2 \circ h = h' \circ g_1$. \mathbb{L}_f^\bullet is then a complex in the derived category $D^-(\mathcal{C})$.

(2) For a B -module M we set

$$T^i(f/A, M) := \text{Ext}_{\mathcal{C}}^i(\mathbb{L}_f^\bullet, (M, M \otimes_B C)), \quad i \geq 0.$$

As before, we simply write $T^i(f, M)$ if $A = \mathbb{C}$ and $T^i(f/A) := T^i(f/A, B)$.

(3) More generally, we write

$$T^i(f/A, (M_1, M_2)) := \text{Ext}_{\mathcal{C}}^i(\mathbb{L}_f^\bullet, (M_1, M_2)), \quad i \geq 0,$$

for any object (M_1, M_2, h) in \mathcal{C} .

Again, one can describe $T^0(f, -)$ more explicitly in terms of derivations:

Proposition 1.6. Let $f : B \rightarrow C$ be as above and let M be a B -module. Then $T^0(f/A, M)$ consists of all pairs of compatible derivations $(\delta, \tilde{\delta}) \in \text{Der}_A(B, M) \times \text{Der}_A(C, M \otimes_B C)$, i. e. pairs $(\delta, \tilde{\delta})$ such that

$$\begin{array}{ccc} C & \xrightarrow{\tilde{\delta}} & M \otimes_B C \\ f \uparrow & & \uparrow \\ B & \xrightarrow{\delta} & M \end{array}$$

commutes.

Furthermore, there is, for any B -module M , a short exact sequence of complexes

$$0 \rightarrow \text{Hom}_C(\Omega_{S/R} \otimes_S C, M \otimes_B C) \rightarrow \text{Hom}_{\mathcal{C}}(\mathbb{L}_f^\bullet, (M, M \otimes_B C)) \rightarrow \text{Hom}_B(\Omega_{R/A} \otimes_R B, M) \rightarrow 0$$

which yields the following long exact cohomology sequence:

Proposition 1.7. For any morphism $f : B \rightarrow C$ of analytic A -algebras and any B -module C there is a long exact sequence in cohomology

$$0 \rightarrow T^0(C/B, M \otimes_B C) \rightarrow T^0(f/A, M) \rightarrow T^0(B/A, M) \rightarrow T^1(C/B, M \otimes_B C) \rightarrow \dots$$

Corollary 1.8. Let $\xi : X \rightarrow S$ be a morphism of complex space germs and M an \mathcal{O}_S -module. Then:

(1) There is a long exact sequence

$$0 \rightarrow T^0(X/S, M \otimes_{\mathcal{O}_S} \mathcal{O}_X) \rightarrow T^0(\xi, M) \rightarrow T^0(S, M) \rightarrow T^1(X/S, M \otimes_{\mathcal{O}_S} \mathcal{O}_X) \rightarrow \dots$$

- (2) Taking $M := \mathcal{O}_S$ one obtains the so-called *Kodaira-Spencer-sequence* for the mapping ξ :

$$0 \rightarrow T^0(X/S) \rightarrow T^0(\xi) \rightarrow T^0(S) \rightarrow T^1(X/S) \rightarrow \dots$$

In particular, for $i = 0$ we have

$$T^0(\xi) = \left\{ (\delta, \tilde{\delta}) \in \text{Der}_{\mathbb{C}}(\mathcal{O}_S, \mathcal{O}_S) \times \text{Der}_{\mathbb{C}}(\mathcal{O}_X, \mathcal{O}_X) : \xi^* \circ \delta = \tilde{\delta} \circ \xi^* \right\},$$

where ξ^* denotes the corresponding homomorphism $\mathcal{O}_S \rightarrow \mathcal{O}_X$.

Again, for small i the modules $T^i f$ have an interpretation in the context of deformations of a morphism $f : X \rightarrow S$: $T^0(f)$ contains the pairs of compatible infinitesimal automorphisms of X and S over the double point D , and $T^1(f)$ coincides with the set of isomorphism classes of deformations of f over D , given by a commutative diagram

$$\begin{array}{ccc} X \hookrightarrow \mathcal{X} & & \\ \downarrow f & \searrow \xi & \\ S \hookrightarrow \mathcal{S} & \nearrow \rho & D, \end{array}$$

$\downarrow F$

where ξ and ρ are deformations of X and S , respectively.

2 Deformations with section

In this section we review the definition of the functor $\text{Def}_{X_0}^s$ of deformations with section of an isolated singularity X_0 and collect some results on the corresponding cotangent cohomology modules and the construction of (mini-)versal deformations with section, see also [Buc81].

2.1 The functor $\text{Def}_{X_0}^s$

Definition. Let X_0 be a germ of a complex space.

- (1) A *deformation with section* of X_0 over a space germ S is a deformation $\xi : X \rightarrow S$ of X_0 together with a section $\sigma : S \rightarrow X$, i. e. $\xi \circ \sigma = \text{id}_S$. That is, we have a cartesian diagram

$$\begin{array}{ccc} 0 \hookrightarrow S & & \\ \downarrow & \searrow \sigma & \\ X_0 \hookrightarrow X & & \\ \downarrow & \searrow \xi & \\ 0 \hookrightarrow S, & & \end{array}$$

$\downarrow i$

and the corresponding homomorphism $\xi^* : \mathcal{O}_S \rightarrow \mathcal{O}_X$ turns \mathcal{O}_X into a flat \mathcal{O}_S -module. We write (ξ, σ) for a deformation together with its section.

- (2) Two deformations $\xi : X \rightarrow S$ and $\xi' : X' \rightarrow S$ over S with sections σ and σ' are *isomorphic* if there is a morphism $\psi : X \rightarrow X'$ preserving X_0 inside X resp. X' and making the diagram

$$\begin{array}{ccc} & S & \\ \sigma \swarrow & & \searrow \sigma' \\ X & \xrightarrow{\psi} & X' \\ \xi \searrow & & \swarrow \xi' \\ & S & \end{array}$$

commute (and ψ is then an isomorphism, whence the terminology).

- (3) *Base change*: If $\varphi : T \rightarrow S$ is a further map of space germs, the *pull-back* (or *induced deformation*) of ξ by φ is defined as the fibre product of X and T over S together with the obvious maps:

$$\begin{array}{ccc} T & \xrightarrow{\varphi} & S \\ \downarrow t \mapsto (\sigma(\varphi(t)), t) & & \downarrow \sigma \\ X \times_S T & \xrightarrow{pr_1} & X \\ \downarrow pr_2 & & \downarrow \xi \\ T & \xrightarrow{\varphi} & S. \end{array}$$

We denote the pull-back as $\varphi^*(\xi)$; if $T \xrightarrow{i} S$ is a subspace we simply write $\xi|_T$ for $i^*(\xi)$.

- (4) With these notions, we can define the *functor of deformations with section of X_0*

$$\begin{aligned} Def_{X_0}^s : (Gan) &\rightarrow (Sets) \\ S &\mapsto \left\{ \begin{array}{l} \text{isomorphism classes of deformations} \\ \text{with section of } X_0 \text{ over } S \end{array} \right\}. \end{aligned}$$

By abuse of notation, we denote by $Def_{X_0}^s$ the (covariant) functor $(Analg) \rightarrow (Sets)$ of deformations with section of the analytic algebra \mathcal{O}_{X_0} , too.

2.2 Cotangent cohomology of $Def_{X_0}^s$

We can interpret deformations with section of X_0 as deformations of the morphism $0 \hookrightarrow X_0$ corresponding to the residue map $\mathcal{O}_{X_0} \twoheadrightarrow \mathbb{C}$. Its cotangent cohomology coincides with the cotangent cohomology of X_0 with values in \mathfrak{m} , the maximal ideal of \mathcal{O}_{X_0} :

Proposition 2.1. Let X_0 be a germ of a complex space, minimally embedded in $(\mathbb{C}^n, 0)$ and given by an ideal $I = (f_1, \dots, f_k) \subseteq \mathcal{O}_n$. Then

$$T^i(0 \hookrightarrow X_0) = T^i(X_0, \mathfrak{m}), i \geq 0,$$

where $\mathfrak{m} \subseteq \mathcal{O}_{X_0}$ is the maximal ideal. In particular we have:

- (1) $T^0(X_0, \mathfrak{m}) \simeq \text{Der}_{\mathbb{C}}(\mathcal{O}_{X_0}, \mathfrak{m})$.

(2) $T^1(X_0, \mathfrak{m})$ is the cokernel of the map

$$\begin{aligned} \mathrm{Der}_{\mathbb{C}}(\mathcal{O}_n, \mathfrak{m}) &\rightarrow \mathrm{Hom}_{\mathcal{O}_{X_0}}(I/I^2, \mathfrak{m}) \\ \delta &\mapsto (g + I^2 \mapsto \delta(g)). \end{aligned}$$

If X_0 is a complete intersection and f_1, \dots, f_k form a regular sequence, then

$$T^1(X_0, \mathfrak{m}) \simeq \mathfrak{m}^{\oplus k} / (\mathfrak{m}J(f)),$$

where $J(f)$ is the module generated by the columns of the Jacobian matrix $\begin{pmatrix} \frac{\partial f_1}{\partial x_1} & \dots & \frac{\partial f_1}{\partial x_n} \\ \vdots & & \vdots \\ \frac{\partial f_k}{\partial x_1} & \dots & \frac{\partial f_k}{\partial x_n} \end{pmatrix}$.

(3) Complete intersections are unobstructed for $Def_{X_0}^s$, i. e. $T^2(X_0, \mathfrak{m}) = 0$.

Proof. Let r denote the residue map $\mathcal{O}_{X_0} \rightarrow \mathbb{C}$. The cotangent complex of \mathbb{C} being trivial we get $\mathrm{Hom}_{\mathcal{C}}(\mathbb{L}_r^\bullet, (\mathcal{O}_{X_0}, \mathbb{C})) \simeq \mathrm{Hom}_{\mathcal{O}_{X_0}}(\mathbb{L}_{\mathcal{O}_{X_0}/\mathbb{C}}^\bullet, \mathfrak{m})$, where \mathcal{C} denotes the category described in section 1.2. Thus

$$\begin{aligned} T^i(0 \hookrightarrow X_0) &= \mathrm{Ext}_{\mathcal{C}}^i(\mathbb{L}_r^\bullet, \mathbb{C}) \\ &\simeq \mathrm{Ext}_{\mathcal{O}_{X_0}}^i(\mathbb{L}_{\mathcal{O}_{X_0}/\mathbb{C}}^\bullet, \mathfrak{m}) = T^i(X_0, \mathfrak{m}). \end{aligned}$$

(1) and (3) are a direct consequence from Proposition 1.1. Property (2) follows from Proposition 1.4, noting that $\mathrm{Der}_{\mathbb{C}}(\mathcal{O}_n, \mathfrak{m})$ is generated by $\left\{x_i \frac{\partial}{\partial x_j}\right\}_{i,j}$ and $\mathrm{Hom}_{\mathcal{O}_{X_0}}((f)/(f^2), \mathfrak{m}) \simeq \mathfrak{m}^{\oplus k}$ for complete intersections. \blacksquare

The same argument as in the case of the functor Def_{X_0} shows:

Lemma 2.2. Let X_0 be a germ of a complex space. Then $T^1(X_0, \mathfrak{m})$ is isomorphic to the \mathcal{O}_{X_0} -module of isomorphism classes of first-order infinitesimal deformations with section of X_0 , where \mathfrak{m} denotes the maximal ideal of \mathcal{O}_{X_0} , i. e. $T^1(X_0, \mathfrak{m})$ coincides with the tangent space of the functor $Def_{X_0}^s$.

Remark. In particular, as embedded deformation, every infinitesimal deformation of $X_0 \subseteq (\mathbb{C}^n, 0)$ given by $f_1, \dots, f_k \in \mathcal{O}_n$ is defined by $f_i + \varepsilon g_i \in \mathcal{O}_n \otimes \mathbb{C}[\varepsilon]/(\varepsilon^2)$, $i = 1, \dots, k$, where the g_i are elements of the maximal ideal of \mathcal{O}_n . More generally, by a suitable coordinate change we may assume that (up to isomorphism) every deformation with section of X_0 over the base S is given by $F_1, \dots, F_k \in (x)\mathcal{O}_S\{x\}$, and σ is the zero section $g \mapsto g \bmod (x)$.

2.3 Construction of a versal deformation with section

We now turn to the construction of a versal deformation with section of an isolated singularity X_0 , which goes back to [Buc81]. There are the usual notions of (mini-)versality:

Definition. Denote by h_Y the $(Sets)$ -valued functor $X \mapsto \mathrm{Hom}(X, Y)$.

- (1) A deformation $\xi : X \rightarrow S$ with section σ is called *versal* (as deformation with section) if for any complex space germ T the map

$$h_S(T) \rightarrow Def_{X_0}^s(T), \quad \varphi \mapsto \varphi^*(\xi)$$

is surjective, i. e. every $(\nu, \tau) \in Def_{X_0}^s(T)$ is induced from ξ by a morphism $T \rightarrow S$.

- (2) As usual, the natural map $\theta_\xi : T(S) \simeq \text{Hom}(\mathcal{O}_S, \mathbb{C}[\varepsilon]) \rightarrow \text{Def}_{X_0}^s(D)$ between the tangent spaces of the functors h_S and $\text{Def}_{X_0}^s$ is called the *Kodaira-Spencer map* of the deformation. (ξ, σ) is called *semi-universal* or *miniversal* (as deformation with section), if it is versal and, in addition, θ_ξ is a bijection.

Theorem 2.3. Suppose $\xi : X \rightarrow S$ is a versal deformation of X_0 . Then $\pi_1 : X \times_S X \rightarrow X$ together with the diagonal embedding $d : X \rightarrow X \times_S X$ gives a versal deformation with section of X_0 .

Proof ([Buc81], see also [MvS01]). Let $\nu : \tilde{X} \rightarrow T$ be a deformation of X_0 with section τ . Versality of ξ implies the existence of some $\varphi : T \rightarrow S$ such that ν is isomorphic to the pull-back of ξ via φ , so we may assume $\tilde{X} = X \times_S T$. Denoting by Φ the projection $\tilde{X} \rightarrow X$ we obtain by pulling back $X \times_S X \rightarrow X$ over $\Phi \circ \tau$:

$$\begin{array}{ccccc}
 & & X \times_S X & \xrightarrow{\pi_2} & X \\
 & \nearrow & \downarrow \pi_1 & & \downarrow \xi \\
 (X \times_S X) \times_X T & & X & \xrightarrow{\xi} & S \\
 \downarrow & \nearrow \Phi \circ \tau & \uparrow \Phi & & \uparrow \varphi \\
 T & \xrightarrow{\tau} & \tilde{X} & \xrightarrow{\nu} & T
 \end{array}$$

From $\tilde{X} = X \times_S T$ we deduce that (up to isomorphism):

$$(X \times_S X) \times_X T \simeq X \times_S (X \times_X T) \simeq X \times_S T = \tilde{X}.$$

Under these identifications, we can identify $d \circ (\Phi \circ \tau)$ with τ . Altogether this proves that (ν, τ) is isomorphic to the pull-back of (π_1, d) via $\Phi \circ \tau$. \blacksquare

Since, by the classical result in [Gra72], any isolated singularity admits a versal deformation, we obtain immediately:

Corollary 2.4. Every isolated singularity X_0 has a (mini-)versal deformation with section.

Example (Complete intersections). We give an explicit construction of a versal deformation of an isolated complete intersection singularity (ICIS) in terms of a defining regular sequence $f = (f_1, \dots, f_k) \in \mathbb{C}\{x_1, \dots, x_n\}^k$: Take a family $\{g^{(1)}, \dots, g^{(\tau)}\}$ of monomials (i. e. $g^{(i)} = x^{\alpha_i} e_{k(i)}$, where e_j denotes the j -th unit vector) representing a \mathbb{C} -vector space basis of $T^1(X_0)$, we may assume that $g^{(i)} = e_i$ for $i = 1, \dots, k$. Then

$$\begin{aligned}
 \mathcal{O}_S &:= \mathbb{C}\{u, s\} / (F_1(u, s), \dots, F_k(u, s)) \\
 &\rightarrow \mathbb{C}\{x, u, s\} / (F_1(x, s), \dots, F_k(x, s), F_1(u, s), \dots, F_k(u, s)) =: \mathcal{O}_X
 \end{aligned}$$

is a versal deformation with section of X_0 , where

$$\begin{pmatrix} F_1(x, s) \\ \vdots \\ F_k(x, s) \end{pmatrix} := \begin{pmatrix} f_1(x) \\ \vdots \\ f_k(x) \end{pmatrix} + \sum_{i=1}^{\tau} s_i g^{(i)}(x)$$

and the section is given by

$$\mathcal{O}_X \rightarrow \mathcal{O}_S, \quad s_i \mapsto s_i, \quad u_i \mapsto u_i, \quad x_i \mapsto u_i.$$

\mathcal{O}_S is regular, using the relation

$$\begin{pmatrix} s_1 \\ \vdots \\ s_k \end{pmatrix} = - \begin{pmatrix} f_1(u) \\ \vdots \\ f_k(u) \end{pmatrix} - \sum_{i=k+1}^{\tau} s_i g^{(i)}(u)$$

in \mathcal{O}_X one obtains that this is isomorphic to

$$\mathbb{C}\{u, s_{k+1}, \dots, s_{\tau}\} \rightarrow \mathbb{C}\{x, u, s_{k+1}, \dots, s_{\tau}\} / (\tilde{F}_1(x, u, s), \dots, \tilde{F}_k(x, u, s)),$$

where

$$\begin{pmatrix} \tilde{F}_1(x, u, s) \\ \vdots \\ \tilde{F}_k(x, u, s) \end{pmatrix} := \begin{pmatrix} f_1(x) - f_1(u) \\ \vdots \\ f_k(x) - f_k(u) \end{pmatrix} + \sum_{i=k+1}^{\tau} s_i (g^{(i)}(x) - g^{(i)}(u)).$$

From the following lemma we can conclude that this deformation is also semi-universal.

At the end of this section a different construction of a semi-universal deformation using a \mathbb{C} -basis of $T^1(X_0, \mathfrak{m})$ is given.

Lemma 2.5. Let $f = (f_1, \dots, f_k) \in \mathcal{O}_n^k$ define an ICIS $X_0 := (V(f), 0)$. Then

$$\tau^s(X_0) := \dim_{\mathbb{C}} T^1(X_0, \mathfrak{m}) = \tau(X_0) + n - k,$$

where $\tau(X_0) =: \dim_{\mathbb{C}} T^1(X_0)$ denotes the Tjurina number of X_0 .

Proof. Denote by (x) the maximal ideal of $\mathcal{O}_n = \mathbb{C}\{x_1, \dots, x_n\}$. By Lemma 2.1 we have

$$T^1(X_0, \mathfrak{m}) = \mathfrak{m}^{\oplus k} / (\mathfrak{m}J(f)) = (x)^{\oplus k} / ((f_i e_j)_{i,j} + (x_i \frac{\partial f}{\partial x_j})_{i,j}).$$

Since $(x)^{\oplus k}$ has \mathbb{C} -codimension k in \mathcal{O}_n^k , the above formula is equivalent to the \mathbb{C} -linear independency of the n partial derivatives $\frac{\partial f}{\partial x_i}$ of f modulo $(f, (x)J(f))$. Suppose the converse, i. e. there exist $\lambda_1, \dots, \lambda_n \in \mathbb{C}$, not all λ_i equal to zero, such that $\sum_i \lambda_i \frac{\partial f}{\partial x_i} \equiv 0 \pmod{(f_i e_j)_{i,j} + (x)J(f)}$. After a linear change of coordinates we may assume that $\frac{\partial f}{\partial x_1} \equiv 0 \pmod{(f_i e_j)_{i,j} + (x)J(f)}$, i. e.

$$\frac{\partial f_i}{\partial x_1} = b f_i + \sum_{j=1}^n c_j \frac{\partial f_i}{\partial x_j}, \quad b \in \mathcal{O}_n, \quad c_j \in (x) \quad (*)$$

for all i . In addition, by a suitable choice of a generating system of the ideal (f_1, \dots, f_k) , we may assume that each generator f_i defines an isolated singularity, hence without loss of generality we may also assume that they are of the form

$$f_i = a_{i1} x_1^{d_{i1}} + \sum_{j=2}^n a_{ij} x_1^{d_{ij}} x_j + r_i(x), \quad a_{ij} \in \mathbb{C}, \quad r_i \in (x_2, \dots, x_n)^2, \quad i = 1, \dots, n,$$

and not all $a_{ij} = 0$, $j = 1, \dots, n$. But taking the partial derivative with respect to x_1 then immediately gives a contradiction to $(*)$. \blacksquare

For arbitrary isolated singularities X_0 defined by $f_1, \dots, f_k \in \mathcal{O}_n$ one may take a different approach to determine (at least k -jets of) a (mini)-versal deformation with section: First compute $g_1, \dots, g_{\tau^s} \in \mathcal{O}_n^k$ representing a \mathbb{C} -vector space basis of $T^1(X_0, \mathfrak{m})$. We obtain a versal family of first order deformations

$$F^{(1)} := f + \sum_{i=1}^{\tau^s} s_i g_i \in (\mathcal{O}_n \otimes \mathbb{C}\{s\}/(s)^2)^k$$

over $\mathbb{C}\{s\}/(s)^2$. Then one lifts this family order by order (in s), killing obstructions. This is the algorithm described in [Mar99] for the computation of versal deformations of singularities, its implementation can be found in the SINGULAR-library `deform.lib` ([Mar02b]).

Since complete intersection are unobstructed, we obtain that in particular a semi-universal deformation with section of an ICIS X_0 is given by

$$F := f + \sum_{i=1}^{\tau^s} s_i g_i \in \mathcal{O}_{n+\tau^s}^k.$$

3 Modular deformations

3.1 Definition and basic properties

A deformation (with section) $\xi : X \rightarrow S$ of an isolated singularity X_0 is called *universal* if every deformation $X' \rightarrow T$ of X_0 is (up to isomorphism) induced by a unique morphism $T \rightarrow S$, i. e. ξ is versal and any two different morphisms $\varphi, \psi : T \rightarrow S$ induce non-isomorphic deformations of X_0 . While any isolated singularity admits a semi-universal deformation (with section), the occurrence of trivial subfamilies in such deformations implies that, in general, universal deformations of X_0 cannot exist, i. e. the functors Def_{X_0} and $Def_{X_0}^s$ are not representable. This is, for instance, the case if X_0 is an isolated complete intersection singularity.

However, restricting a semi-universal family to subgerms of the base for which the universality condition holds is a possible approach to the construction of local moduli for singularities. We call such subgerms *modular*, as introduced by Palamodov in [Pal78]. The similar notion of *prorepresenting substratum* inside the base space of a semi-universal deformation of X_0 is – in the context of formal deformations – studied in [LP88].

Definition. Let $\xi : X \rightarrow S$ be a deformation of X_0 (with section σ). A subspace $M \subseteq S$ is called *modular* if the following condition holds: If $\varphi : T \rightarrow M$ and $\psi : T \rightarrow S$ are morphisms such that the induced deformations $\varphi^*(\xi|_M)$ and $\psi^*(\xi)$ with base T are isomorphic as deformations (with section) then $\varphi = \psi$.

The restriction of a deformation to a modular subgerm is called a *modular deformation*.

Remark. The following properties are immediate from the definition:

- (1) Any subgerm of a modular germ is again modular.
- (2) $\{0\} \subseteq S$ is a modular subspace for the deformation $\xi : X \rightarrow S$ (with section σ) if and only if the corresponding Kodaira-Spencer map $\theta_\xi : T(S) \rightarrow T^1(X_0)$ (resp. $\theta_\xi : T(S) \rightarrow T^1(X_0, \mathfrak{m})$) is injective. Together with (1) this implies that S contains a modular subspace if and only if the Kodaira-Spencer map of the deformation is injective. Following [Pal90] we call such deformations *monodeformations*.

- (3) If M_1, M_2 are modular subgerms of S , then $M_1 \cup M_2 \subseteq S$ is modular, too.
- (4) As a consequence of the identity theorem for power series, two morphisms between space germs $\varphi, \psi : T \rightarrow S$ coincide if $\varphi|_W = \psi|_W$ for any Artinian subgerm W of T , thus it suffices to check the condition of the definition for Artinian germs T .
- (5) There is a unique isomorphism between any two maximal modular subgerms in the base space of a semi-universal deformation of X_0 . Such a maximal modular subgerm is then called the *modular stratum* of X_0 .

3.2 The Kodaira-Spencer sequence of a deformation

In order to derive a criterion for $M \subseteq S$ to be modular, we consider the following commutative diagram with exact rows, combining the Kodaira-Spencer sequence for $\xi : X \rightarrow S$ (Corollary 1.8 (2)) and its evaluation at the special fibre over $0 \in S$:

$$\begin{array}{ccccccc}
0 & \longrightarrow & T^0(X/S) & \longrightarrow & T^0(\xi) & \longrightarrow & T^0(S) \xrightarrow{\Theta_\xi} T^1(X/S) \longrightarrow \cdots \\
& & \downarrow ev & & \downarrow ev' & & \downarrow \\
0 & \longrightarrow & T^0(X_0) & \longrightarrow & T^0(\xi, (\mathbb{C}, \mathcal{O}_{X_0})) & \longrightarrow & T^0(S, \mathbb{C}) \xrightarrow{\theta_\xi} T^1(X_0) \longrightarrow \cdots,
\end{array} \quad (3.1)$$

cf. [Pal90, § 1]. Evaluation at the special fibre means taking cohomology with values in $\mathbb{C} \simeq \mathcal{O}_S/\mathfrak{m}_S$ instead of \mathcal{O}_S . Furthermore in the lower row we apply Proposition 1.5. to obtain $T^i(X/S, M) \simeq T^i(X_0, M)$ for any $i \geq 0$ and any \mathcal{O}_{X_0} -module M . Here θ_ξ denotes the *Kodaira-Spencer map* of the deformation, and Θ_ξ is the corresponding *relative Kodaira-Spencer map* $T^0(S) \rightarrow T^1(X/S)$.

The following proposition gives an analogue of sequence (3.1) for deformations with section.

Proposition 3.1. Let $\xi : X \rightarrow S$ be a deformation of X_0 with section σ , $J_\sigma \subseteq \mathcal{O}_X$ the kernel of σ^* . Then the sequence

$$0 \longrightarrow T^0(X/S, J_\sigma) \longrightarrow T^0(\xi, (\mathcal{O}_S, J_\sigma)) \longrightarrow T^0(S) \xrightarrow{\Theta_\xi} T^1(X/S, J_\sigma) \longrightarrow \cdots \quad (3.2)$$

is exact and by evaluation at the special fibre over $0 \in S$ we obtain the diagram with exact rows

$$\begin{array}{ccccccc}
0 & \longrightarrow & T^0(X/S, J_\sigma) & \longrightarrow & T^0(\xi, (\mathcal{O}_S, J_\sigma)) & \longrightarrow & T^0(S) \xrightarrow{\Theta_\xi} T^1(X/S, J_\sigma) \longrightarrow \cdots \\
& & \downarrow ev & & \downarrow ev' & & \downarrow \\
0 & \longrightarrow & T^0(X_0, \mathfrak{m}) & \longrightarrow & T^0(\xi, (\mathbb{C}, \mathfrak{m})) & \longrightarrow & T^0(S, \mathbb{C}) \xrightarrow{\theta_\xi} T^1(X_0, \mathfrak{m}) \longrightarrow \cdots,
\end{array} \quad (3.3)$$

where \mathfrak{m} denotes the maximal ideal of \mathcal{O}_{X_0} . We call (3.2) the *Kodaira-Spencer sequence* and Θ_ξ the *relative Kodaira-Spencer map* of (ξ, σ) .

Proof. Choose resolvents R_X and R_S of \mathcal{O}_X resp. \mathcal{O}_S over \mathbb{C} as in the definition of the cotangent complex $\mathbb{L}_{\xi^*}^\bullet$ in section 1.2. The section σ^* yields a split exact sequence

$$0 \longrightarrow J_\sigma \xrightleftharpoons[\tau]{\iota} \mathcal{O}_X \xrightleftharpoons[\xi^*]{\sigma^*} \mathcal{O}_S \longrightarrow 0,$$

hence we obtain a homomorphism $\tau \circ \xi^* : \mathcal{O}_S \rightarrow J_\sigma$ over ξ^* , i. e. $(\mathcal{O}_S, J_\sigma, \tau \circ \xi^*)$ is an object in the category $\mathcal{C}_{\xi^* : \mathcal{O}_S \rightarrow \mathcal{O}_X}$ defined in section 1. All cotangent complexes fit together into the short exact sequence

$$0 \rightarrow \mathrm{Hom}_{\mathcal{O}_X}(\Omega_{R_X/R_S} \otimes_{R_X} \mathcal{O}_X, J_\sigma) \rightarrow \mathrm{Hom}_{\mathcal{C}}(\mathbb{L}_{\xi^*}^\bullet, (\mathcal{O}_S, J_\sigma, \tau \circ \xi^*)) \rightarrow \mathrm{Hom}_{\mathcal{O}_S}(\mathbb{L}_{\mathcal{O}_S/\mathbb{C}}^\bullet, \mathcal{O}_S) \rightarrow 0.$$

The choice of R_X and R_S implies that $\Omega_{R_X/R_S} \otimes_{R_X} \mathcal{O}_X \simeq \mathbb{L}_{\mathcal{O}_X/\mathcal{O}_S}^\bullet$ (cf. [Fle78, ch. I]), hence we get (3.2) as its long exact cohomology sequence.

Similarly, if one takes coefficients in $(\mathbb{C}, \mathfrak{m})$, this produces the exact lower row of (3.3), and the commutativity of this diagram is then clear from functoriality. \blacksquare

3.3 A criterion for modularity in terms of T^0

We are now going to give a criterion for $M \subseteq S$ to be modular, similar statements for deformations of complex spaces are already contained in [Pal78]. In [KS90] modular subspaces for general deformation groupoids are studied, and an analogue for criterion (3) below is given, involving a so-called *exponential functor* that generalizes the module of derivations $T^0(X/S)$. We use the following two auxiliary notions, the first of which being motivated by the fact that $\{0\} \subseteq S$ is modular if and only if θ_ξ is injective.

Definition. A subgerm $M \subseteq S$ in the base space of a deformation $\xi : X \rightarrow S$ (with section σ) of X_0 is called

- (1) *infinitesimally modular* if the restriction to M of the map

$$\tilde{\Theta}_{\xi|M} : T^0(S, \mathcal{O}_M) \rightarrow T^1(X/S)|_M$$

(resp. $\tilde{\Theta}_{\xi|M} : T^0(S, \mathcal{O}_M) \rightarrow T^1(X/S, J_\sigma)|_M$ if ξ is considered as deformation with section σ) is injective;

- (2) *Artinian modular* if any Artinian subgerm of M is modular.

(Here and subsequently we use the notations $T^i(X/S)|_M := T^i(X|_M/M)$ etc. for the cotangent cohomology of the restricted deformation $\xi|_M$.)

Lemma 3.2. For both deformation functors, modular subspaces are infinitesimally modular.

Proof (cf. [Mar02a]). Suppose $\delta \in \mathrm{Ker}(\tilde{\Theta}_{\xi|M}) \subseteq T^0(S, \mathcal{O}_M)$. Then δ corresponds to a morphism $\varphi_\delta : M \times D \rightarrow S$ such that the deformation $\varphi_\delta^*(\xi)$ is isomorphic to the trivial one, which can also be induced from $\xi|_M$ via the projection $pr_M : M \times D \rightarrow M$. By the modularity of M we conclude that $\varphi_\delta = i \circ pr_M$, hence $\delta = 0$. \blacksquare

Lemma 3.3. Suppose $\xi : X \rightarrow S$ is either a semi-universal or a modular deformation of X_0 (with section σ). Then, in each case, the mapping $ev' : T^0(\xi) \rightarrow T^0(\xi, (\mathbb{C}, \mathcal{O}_{X_0}))$ (resp. $ev' : T^0(\xi, (\mathcal{O}_S, J_\sigma)) \rightarrow T^0(\xi, (\mathbb{C}, \mathfrak{m}))$) is surjective.

Proof. First suppose ξ is semi-universal. Take a derivation $\delta \in T^0(\xi, (\mathbb{C}, \mathfrak{m}))$. Since ξ is semi-universal, we may interpret δ as element of $T^0(X_0, \mathfrak{m})$ and it suffices to find a preimage in $T^0(\xi, (\mathcal{O}_S, J_\sigma))$.

We use the construction of [Pal90, Proposition 18.9]: δ induces an automorphism a_δ of $X_0 \times D$ over D . Denote by $\tilde{\xi}$ the deformation $\xi \times id_D : X \times D \rightarrow S \times D$ with section $\sigma \times id_D$. Let $T := (\{0\} \times D) \cup (S \times \{0\})$ and $\varphi : T \rightarrow S$ be the canonical projection onto S , so $\tilde{\xi}|_T = \varphi^*(\xi)$. Let \tilde{a}_δ be the automorphism of $\varphi^*(\xi)$ induced by a_δ on the first component of T and by id_S on the second. Then $\xi' := \tilde{a}_\delta \circ \varphi^*(\xi)$ is of course still a versal deformation with section (isomorphic to $\tilde{\xi}|_T$), hence $\tilde{\xi}$ is induced from it by some $\psi : S \times D \rightarrow X$, which altogether yields the diagram

$$\begin{array}{ccccc} a' : X \times D & \longrightarrow & (X \times D)|_T & \xrightarrow{pr_X \circ \tilde{a}_\delta} & X \\ \downarrow \tilde{\xi} & & \downarrow \xi' & & \downarrow \xi \\ \psi' : S \times D & \xrightarrow{\psi} & T & \xrightarrow{\varphi} & S \end{array}$$

The corresponding map $\psi'^* : \mathcal{O}_S \rightarrow \mathcal{O}_S[\varepsilon]$ gives a vector field $\eta \in \text{Der}_{\mathbb{C}}(\mathcal{O}_S)$ defined by $\psi'^*(a) = a + \varepsilon\eta(a)$, similarly a' induces a vector field $\tilde{\delta} \in \text{Der}_{\mathbb{C}}(\mathcal{O}_X, J_\sigma)$, with the property that $(\tilde{\delta}, \eta) \in T^0(\xi, (\mathcal{O}_S, J_\sigma))$ and $ev'(\tilde{\delta}, \eta) = \delta$.

Now let $\hat{\xi} : Y \rightarrow M$ be a modular deformation with section $\hat{\sigma}$. We may assume that M is a subspace of the base of a semi-universal deformation with section $\xi : X \rightarrow S$ of X_0 , and $\hat{\xi} = \xi|_M$. As before, let $\tilde{\xi} := \hat{\xi} \times id_D$, then $\delta \in T^0(\hat{\xi}, (\mathbb{C}, \mathfrak{m})) \simeq T^0(X_0, \mathfrak{m})$ induces an automorphism \tilde{a}_δ of the (still versal) deformation $\tilde{\xi}|_T$, so we can induce $\tilde{\xi} : Y \times D \rightarrow M \times D$ from $\xi' := a_\delta \circ \tilde{\xi}|_T$ by means of some morphism ψ :

$$\begin{array}{ccccc} a' : Y \times D & \longrightarrow & (X \times D)|_T & \xrightarrow{pr_X \circ \tilde{a}_\delta} & X \\ \downarrow \tilde{\xi} & & \downarrow \xi' & & \downarrow \xi \\ \psi' : M \times D & \xrightarrow{\psi} & T & \xrightarrow{\varphi = pr_S} & S \end{array}$$

On the other hand we can obtain $\tilde{\xi}$ from ξ by just taking the projection $pr_1 : M \times D \rightarrow M \subseteq S$. Thus, by the modularity of M , $pr_1 = \psi'$, so that a' and ψ' can be considered as maps $Y \times D \rightarrow Y$ resp. $M \times D \rightarrow M$, inducing the desired lift of δ to $\text{Der}_{\mathbb{C}}(\mathcal{O}_Y, J_{\hat{\sigma}})$ resp. $\text{Der}_{\mathbb{C}}(\mathcal{O}_M)$. \blacksquare

With these preparations we can formulate and prove the following criterion for modularity:

Theorem 3.4. For a subspace $M \subseteq S$ of the base space of a monodeformation $\xi : X \rightarrow S$ of X_0 the following assertions are equivalent:

- (1) M is modular.
- (2) M is infinitesimally modular.
- (3) $ev|_M : T^0(X/S)|_M \rightarrow T^0(X_0)$ is surjective.
- (4) M is Artinian modular.

Theorem 3.5. For a subspace $M \subseteq S$ of the base space of a monodeformation $\xi : X \rightarrow S$ of X_0 with section $\sigma : S \rightarrow X$ the following assertions are equivalent:

- (1) M is modular.
- (2) M is infinitesimally modular.
- (3) $ev|_M : T^0(X/S, J_\sigma)|_M \rightarrow T^0(X_0, \mathfrak{m})$ is surjective.
- (4) M is Artinian modular.

We only give a proof of Theorem 3.5 – Theorem 3.4 follows by omitting any section occurring and replacing any $T^i(X_0, \mathfrak{m})$ by $T^i(X_0)$, $T^i(X/S, J_\sigma)$ by $T^i(X/S)$ etc.

Proof (of Theorem 3.5). (1) \Rightarrow (2) : This is Lemma 3.2.

(2) \Rightarrow (3) : Consider the Kodaira-Spencer sequence (3.2) of (ξ, σ) with coefficients in $(\mathcal{O}_M, J_{\sigma|_M})$. This sequence can be put in between the rows of (3.3) so that, altogether, we obtain the commutative diagram with exact rows

$$\begin{array}{ccccccc}
0 & \longrightarrow & T^0(X/S, J_\sigma) & \longrightarrow & T^0(\xi, (\mathcal{O}_S, J_\sigma)) & \longrightarrow & T^0(S) \xrightarrow{\Theta_\xi} T^1(X/S, J_\sigma) \longrightarrow \dots \\
& & \tilde{ev} \downarrow & & \tilde{ev}' \downarrow & & \downarrow \\
0 & \longrightarrow & T^0(X/S, J_\sigma)|_M & \longrightarrow & T^0(\xi, (\mathcal{O}_M, J_{\sigma|_M})) & \longrightarrow & T^0(S, \mathcal{O}_M) \xrightarrow{\tilde{\Theta}_{\xi|_M}} T^1(X/S, J_\sigma)|_M \longrightarrow \dots \\
& & ev|_M \downarrow & & ev'|_M \downarrow & & \downarrow \\
0 & \longrightarrow & T^0(X_0, \mathfrak{m}) & \longrightarrow & T^0(\xi, (\mathbb{C}, \mathfrak{m})) & \longrightarrow & T^0(S, \mathbb{C}) \xrightarrow{\theta_\xi} T^1(X_0, \mathfrak{m}) \longrightarrow \dots
\end{array} \tag{3.4}$$

By assumption $\tilde{\Theta}_{\xi|_M}$ is injective, $\theta_{\xi|_M}$ is injective since ξ is a monodeformation. Thus in (3.4) we can identify $ev|_M$ and $ev'|_M$, the latter being surjective since, by Lemma 3.3, the composition $ev' = ev'|_M \circ \tilde{ev}'$ is surjective.

(3) \Rightarrow (4) : Suppose $ev|_M$ is surjective and take an Artinian subgerm $A \subseteq M$. Since A is obtained from \mathbb{C} by a finite number of small extensions and $\{0\} \subseteq S$ is modular if and only if θ_ξ is injective, it suffices to prove the following statement: Let ξ be modular over the Artinian subgerm A_0 of M , then it is also modular over A given by an infinitesimal extension

$$\eta : 0 \rightarrow K \rightarrow \mathcal{O}_A \rightarrow \mathcal{O}_{A_0} \rightarrow 0.$$

As in [Mar, Proposition 1.4] one associates to η the obstruction element

$$ob_{\xi, \eta} \in \text{Hom}(T^0(X_0, \mathfrak{m}), T^1(X_0, \mathfrak{m})) \otimes K,$$

whose vanishing is then equivalent to A being modular, which is proved absolutely analogously to [Mar, Lemma 1.5]. In concrete terms, we may assume that $\mathcal{O}_A = \mathcal{O}_{A_0}[\varepsilon]$ and $K = (\varepsilon)$, $\varepsilon^2 = 0$. Furthermore let X_0 be given by $f = (f_1, \dots, f_k) \in \mathcal{O}_n^k$, X be defined by $F = (F_1, \dots, F_k) \in (\mathcal{O}_n \otimes \mathcal{O}_S)^k$.

Then we can write $F|_A = F|_{A_0} + \varepsilon \tilde{f}$ with $\varepsilon \tilde{f} \in \mathcal{O}_n^k$ defining a class in $T^1(X_0, \mathfrak{m}) \otimes K$. Any $\delta \in T^0(X_0, \mathfrak{m})$ is represented by $\delta' \in \text{Der}_{\mathbb{C}}(\mathcal{O}_n, (x))$ with $\delta'(f) = h \cdot f$ for some matrix h with entries in \mathcal{O}_n . By assumption (3) we can lift δ to some $\delta|_{A_0} \in T^0(X|_{A_0}, J_{\sigma|_{A_0}})$, represented by

$\delta'_{|A_0}$ with $\delta'_{|A_0}(F_{|A_0}) = H_{|A_0}F_{|A_0}$, $H_{|A_0}$ being a lift of h to $\mathcal{O}_n \otimes \mathcal{O}_{A_0}$. A lift of $\delta'_{|A_0}$ and $H_{|A_0}$ to $\delta'_{|A}$ and $H_{|A}$ over A induces classes

$$\delta'_{|A}(F_{|A}) - H_{|A}F_{|A} = (\delta'(f) - h\tilde{f})\varepsilon \in T^1(X_0, \mathfrak{m}) \otimes K,$$

which yields a homomorphism $ob_{\xi, \eta} : T^0(X_0, \mathfrak{m}) \rightarrow T^1(X_0, \mathfrak{m}) \otimes K$. In fact, ob is induced by the Lie bracket $[-, -] : T^0(X_0, \mathfrak{m}) \times T^1(X_0, \mathfrak{m}) \rightarrow T^1(X_0, \mathfrak{m})$ which implies that it does not depend on the choices made.

Now by assumption (3) there exists a lift $\delta'_{|A} \in T^0(X_{|A}, J_{\sigma|A})$, thus $ob_{\xi, \eta}$ vanishes, hence A is modular.

(4) \Rightarrow (1): Let Z be an Artinian germ and $\varphi : Z \rightarrow M$, $\psi : Z \rightarrow S$ morphisms such that $\varphi^*(\xi_{|M}) \simeq \psi^*(\xi)$. We can factor $\varphi = i \circ \varphi_0$, where $\varphi_0 : Z \rightarrow M_0$ with M_0 Artinian and $i : M_0 \hookrightarrow M$ the inclusion. Thus $\psi^*(\xi) \simeq \varphi_0^*(i^*(\xi_{|M})) \simeq \varphi_0^*(\xi_{|M_0})$. By the assumption M_0 is modular, so $\psi = \varphi_0$ and therefore ψ and φ coincide, too. \blacksquare

Remark. The definition of infinitesimal modularity used here differs slightly from the one used in [Mar02a], [Mar], where it is characterized as injectivity of the Kodaira-Spencer map $\Theta_{\xi|_M} : T^0(M) \rightarrow T^1(X/S)_{|M}$ of the deformation $\xi_{|M}$. For any subspace $M \subseteq S$ we have a commutative diagram

$$\begin{array}{ccc} T^0(S, \mathcal{O}_M) & \xrightarrow{\tilde{\Theta}_{\xi|_M}} & T^1(X_{|M}/M), \\ \uparrow & \nearrow \Theta_{\xi|_M} & \\ T^0(M) & & \end{array}$$

so $\text{Ker}(\tilde{\Theta}_{\xi|_M}) = 0$ is the stronger notion, and it is indeed this condition that we need in the above proof of the implication (2) \Rightarrow (3).

As a corollary of the obstruction calculus in the proof of implication (3) \Rightarrow (4) we obtain:

Corollary 3.6. The tangent space $T(M)$ to the modular stratum of X_0 equals the subspace

$$\{t \in T(S) : [\theta_{\xi}(t), \delta] = 0 \text{ for all } \delta \in T^0(X_0)\}$$

(resp. $T^0(X_0, \mathfrak{m})$ for deformations with section).

3.4 The module $T^0_{\bullet}(X_0)$

Obviously, surjectivity of the evaluation mappings in the above statements is equivalent to the surjectivity of the mapping

$$T^0(X/S)_{|M} \rightarrow T^0(X_0)/\text{ev}(T^0(X/S)) \quad \text{resp.} \quad T^0(X/S, J_{\sigma})_{|M} \rightarrow T^0(X_0, \mathfrak{m})/\text{ev}(T^0(X/S, J_{\sigma})).$$

This gives rise to the following notions (cf. [Pal90]):

Definition. Let $\xi : X \rightarrow S$ be a semi-universal deformation of X_0 (with section σ). We set

- $T^0_{\bullet}(X_0) := T^0(X_0)/\text{ev}(T^0(X/S))$, resp.
- $T^0_{\bullet}(X_0, \mathfrak{m}) := T^0(X_0, \mathfrak{m})/\text{ev}(T^0(X/S, J_{\sigma}))$ for the case of deformations with section.

We note some properties of $T^\bullet_\bullet(X_0)$ stated in [Pal90]:

- (1) $\dim_{\mathbb{C}} T^\bullet_\bullet(X_0) < \infty$;
- (2) $\dim_{\mathbb{C}} T^\bullet_\bullet(X_0) \leq \tau(X_0)$, with equality if and only if the kernel of the Kodaria-Spencer map $T^0(S) \rightarrow T^1(X/S)$ is free. This is, for instance, the case if X_0 is a complete intersection (by the Saito-Looijenga-Theorem, see [Sai80], [Loo84]) or if X_0 is a reduced space curve singularity, as was shown by D. van Straten in [vS95];
- (3) if X_0 is a hypersurface defined by $f \in \mathcal{O}_n$, then there is an exact sequence of \mathcal{O}_n -modules

$$0 \rightarrow T^\bullet_\bullet(X_0) \xrightarrow{\varepsilon} \mathcal{O}_n/J(f) \xrightarrow{m_f} \mathcal{O}_n/J(f) \xrightarrow{\pi} T^1(X_0) \rightarrow 0, \quad (3.5)$$

where π is the projection, m_f denotes multiplication by f modulo $J(f)$ and ε is defined as follows: If $\delta \in T^0(X_0)$ is the class of a derivation $\delta' : \mathcal{O}_n \rightarrow \mathcal{O}_n$ subject to $\delta'(f) = h \cdot f$ for some $h \in \mathcal{O}_n$, then define $\varepsilon(\delta) := \frac{\delta'(f)}{f} \bmod J(f)$.

Properties similar to (1) and (3) above hold for the module $T^\bullet_\bullet(X_0, \mathfrak{m})$.

Proposition 3.7. Let $\xi : X \rightarrow S$ be a semi-universal deformation of X_0 with section σ . Then:

- (1) $\dim_{\mathbb{C}} T^\bullet_\bullet(X_0, \mathfrak{m}) < \infty$;
- (2) if X_0 is a hypersurface defined by $f \in \mathcal{O}_n$, then there is an exact sequence of \mathcal{O}_n -modules

$$0 \rightarrow T^\bullet_\bullet(X_0, \mathfrak{m}) \xrightarrow{\varepsilon} \mathcal{O}_n/J(f) \xrightarrow{m_f} \mathcal{O}_n/(x)J(f) \xrightarrow{\pi} \mathcal{O}_n/(f, (x)J(f)) \rightarrow 0, \quad (3.6)$$

where the maps ε, m_f and π are defined analogously as in (3.5). In particular we have:

$$\dim_{\mathbb{C}} T^\bullet_\bullet(X_0, \mathfrak{m}) = \dim_{\mathbb{C}} T^\bullet_\bullet(X_0) = \tau.$$

Proof. Property (1) is a consequence of the Kodaira-Spencer sequence (Proposition 3.1), cf. [Pal90, Proposition 1.7]: ξ being semi-universal, θ_ξ is injective, thus we may again identify $T^0(X_0, \mathfrak{m})$ and $T^0(\xi, (\mathbb{C}, \mathfrak{m}))$. Together with the surjectivity of the map ev' (proved in Lemma 3.3) we obtain that ev' induces a surjective \mathcal{O}_S -linear map

$$T^0(\xi, (\mathcal{O}_S, J_\sigma))/T^0(X/S, J_\sigma) \rightarrow T^\bullet_\bullet(X_0, \mathfrak{m}).$$

The \mathcal{O}_S -module on the left is isomorphic to the submodule $Im(T^0(\xi, (\mathcal{O}_S, J_\sigma)) \rightarrow T^0(S)) = Ker(\Theta_\xi)$ of the finitely generated \mathcal{O}_S -module $T^0(S)$, hence it is itself finitely generated. Now applying $-\otimes_{\mathcal{O}_S} \mathbb{C}$ yields a surjective map

$$T^0(\xi, (\mathcal{O}_S, J_\sigma))/T^0(X/S, J_\sigma) \otimes_{\mathcal{O}_S} \mathbb{C} \rightarrow T^\bullet_\bullet(X_0, \mathfrak{m}) \otimes_{\mathcal{O}_S} \mathbb{C} \simeq T^\bullet_\bullet(X_0, \mathfrak{m}),$$

whose source is a finite-dimensional \mathbb{C} -vector space, hence $\dim_{\mathbb{C}} T^\bullet_\bullet(X_0, \mathfrak{m}) < \infty$, too.

We now prove (2), which is done analogously to [Pal90, Proposition 3.1]. ε is well-defined: Let X be defined by $F \in \mathcal{O}_S\{x\}$, in addition we may assume that σ is the zero section, i. e. $J_\sigma = (x)\mathcal{O}_S\{x\}$. If $\delta \in T^0(X_0, \mathfrak{m})$ lifts to $\tilde{\delta} \in T^0(X/S, J_\sigma)$, then there exist $G_i \in (x)\mathcal{O}_S\{x\}$ and $H \in \mathcal{O}_S\{x\}$ such that δ is the class of $\sum_i G_i \frac{\partial}{\partial x_i}$ and $\sum_i G_i \frac{\partial F}{\partial x_i} = H \cdot F$. Since F and its

partial derivatives with respect to the x_i form a regular sequence in $\mathcal{O}_S\{x\}$, this implies that H is in the ideal generated by $\frac{\partial F}{\partial x_1}, \dots, \frac{\partial F}{\partial x_n}$, i. e. $h = H|_{s=0} \in J(f)$.

It remains to verify the exactness of the sequence, which is obvious at the two terms on the right. If $h \bmod J(f) \in \text{Ker}(m_f)$, then $hf = \sum_i g_i \frac{\partial f}{\partial x_i}$ with $g_i \in (x)$, i. e. $h = \varepsilon(\delta)$ for the derivation $\delta : \mathcal{O}_{X_0} \rightarrow \mathfrak{m}$ induced by $\sum_i g_i \frac{\partial f}{\partial x_i}$. Since the inclusion $\text{Im}(\varepsilon) \subseteq \text{Ker}(m_f)$ is clear, it is only left to show that ε has trivial kernel. But if δ is induced by a derivation δ' of \mathcal{O}_n with $\sum_i g_i \frac{\partial f}{\partial x_i} = h \cdot f$ and $h = \sum_i h_i \frac{\partial f}{\partial x_i}$, then its class equals the class of $\sum_i (g_i - f \cdot h_i) \frac{\partial f}{\partial x_i}$, so we may assume that for its lift δ' we have $\delta'(f) = 0$. Thus its coefficients are a syzygy of the partial derivatives of f which form a regular sequence in \mathcal{O}_n . Hence δ belongs to the submodule generated by the classes of the derivations of the form $\frac{\partial f}{\partial x_j} \frac{\partial}{\partial x_i} - \frac{\partial f}{\partial x_i} \frac{\partial}{\partial x_j}$, which clearly can be lifted to $T^0(X/S)$.

The last claim is now a consequence of Lemma 2.5, which implies that

$$\dim_C \mathcal{O}_n/(f, (x)J(f)) = \dim_C T^1(X_0, \mathfrak{m}) + 1 = \tau(X_0) + n.$$

Now let, in addition, $\mu(X_0) := \dim_C \mathcal{O}_n/J(f)$ be the Milnor number of X_0 . From the fact that the partial derivatives of f form a regular sequence, we conclude that they are linearly independent modulo $(x)J(f)$, hence $\dim_C \mathcal{O}_n/(x)J(f) = \mu + n$. Altogether we calculate using (3.5):

$$\begin{aligned} \dim_C T_\bullet^0(X_0) &= \dim_C \mathcal{O}_n/J(f) - \dim_C \mathcal{O}_n/(x)J(f) + \dim_C \mathcal{O}_n(f, (x)J(f)) \\ &= \mu - (\mu + n) + (\tau(X_0) + n) \\ &= \tau(X_0), \end{aligned}$$

which proves the last assertion. ■

3.5 Quasihomogeneous complete intersections

In general, the modular stratum of an isolated singularity X_0 carries a non-reduced structure. This is, for instance, the case for semi-quasihomogeneous hypersurface singularities. However, it is proved in [Ale85, section 6.2], that the modular stratum is reduced and smooth in case X_0 is a quasihomogeneous isolated complete intersection singularity. As an application of the criteria for modularity established above, we extend this result to deformations with section:

Proposition 3.8. The modular stratum with respect to the functor $\text{Def}_{X_0}^s$ of a quasihomogeneous isolated complete intersection singularity X_0 is reduced and smooth.

Proof. Let X_0 be defined by quasihomogeneous polynomials $f = (f_1, \dots, f_k) \in \mathcal{O}_n^k$ of degrees d_1, \dots, d_k with respect to some positive integer weights w_1, \dots, w_n . For any $1 \leq i_1 < \dots < i_{k+1} \leq n$ one obtains a *Hamiltonian derivation* by cofactor expansion with respect to the first line of the symbolic matrix

$$H(i_1, \dots, i_{k+1}) := \det \begin{vmatrix} \frac{\partial}{\partial x_{i_1}} & \dots & \frac{\partial}{\partial x_{i_{k+1}}} \\ \frac{\partial f_1}{\partial x_{i_1}} & \dots & \frac{\partial f_1}{\partial x_{i_{k+1}}} \\ \vdots & & \vdots \\ \frac{\partial f_k}{\partial x_{i_1}} & \dots & \frac{\partial f_k}{\partial x_{i_{k+1}}} \end{vmatrix},$$

and it is proved in [Ale85, section 6.1] that the Hamiltonian derivations together with the Euler derivation $\delta_E := \sum_{i=1}^n w_i x_i \frac{\partial}{\partial x_i}$ generate the \mathcal{O}_{X_0} -module $T^0(X_0) = \text{Der}_{\mathbb{C}}(\mathcal{O}_{X_0})$. Note that in particular all these derivations take their image in the maximal ideal \mathfrak{m} of \mathcal{O}_{X_0} , so in this situation we have $T^0(X_0) = T^0(X_0, \mathfrak{m})$ and we can take them as generators of the latter module, too.

Now let $\xi : X \rightarrow S$ be a semi-universal deformation with section σ of X_0 . In particular, by the results of section 2.3, $S \simeq (\mathbb{C}^s, 0)$ is smooth. Obviously, all $H(i_1, \dots, i_{k+1})$ can be lifted to $T^0(X/S, J_\sigma)$, so we only have to consider δ_E . The bracket $[-, -]$ in cotangent cohomology induces an element $[\delta_E, -] \in \text{End}_{\mathbb{C}}(T^1(X_0, \mathfrak{m}))$ that gives a decomposition $T^1(X_0, \mathfrak{m}) = \bigoplus_{\nu \in \mathbb{Z}} T^1(X_0, \mathfrak{m})_\nu$ into a direct sum of its eigenspaces $T^1(X_0, \mathfrak{m})_\nu$ with respect to the eigenvalue ν . Using the description of the tangent space to the modular stratum M in Corollary 3.6 we obtain $T(M) \simeq T^1(X_0, \mathfrak{m})_0$.

But now it is clear that there are no further obstructions to lifting δ_E : If $F = f + \sum_{i=1}^{\tau^s} s_i g^{(i)}$ is a semi-universal family with monomials $g^{(i)}$ as in section 2.3 above, then we may assume that $g^{(1)}, \dots, g^{(r)}$ represent a basis of $T^1(X_0, \mathfrak{m})_0$. From what was said above, it then follows that $T(M) \simeq ((s_1, \dots, s_r)/(s_1, \dots, s_r)^2)^*$, and we can lift δ_E to the restriction of ξ to the smooth subspace $(\mathbb{C}^r \times \{0\}, 0)$ of S , i. e. $\xi|_{(\mathbb{C}^r \times \{0\}, 0)}$ is the maximal modular deformation inside ξ . \blacksquare

The argument given above is entirely the same in the context of deformation without section, it is a consequence of the identity $T^0(X_0) = T^0(X_0, \mathfrak{m})$ for singularities of this type. As a corollary we obtain:

Corollary 3.9. For any quasihomogeneous isolated complete intersection singularity X_0 the modular strata with respect to both deformation functors Def_{X_0} and $\text{Def}_{X_0}^s$ coincide.

4 Flatness and modular deformations

4.1 Modularity as flatness of the first Tjurina module

Now choose a minimal embedding $(X_0, 0) \subseteq (\mathbb{C}^n, 0)$. Let X_0 be defined by $f_1, \dots, f_k \in \mathcal{O}_n$ and let $\xi : X \rightarrow S$ be a deformation of X_0 , given by $F_1, \dots, F_k \in \mathcal{O}_S\{x\}$. Denote by $J(f)$ the Jacobian matrix of f_1, \dots, f_k and by $J(F)$ the relative (with respect to x_1, \dots, x_n) Jacobian matrix of F_1, \dots, F_k . Finally, set $\tilde{T}^1(X/S) := \mathcal{O}_X^k / J(F)$. With these notions we can interpret the criterion in Theorem 3.4 as flatness of the module $\tilde{T}^1(X/S)$:

Proposition 4.1. Let $M \subseteq S$ be a subspace in the base of a semi-universal deformation $\xi : X \rightarrow S$ of unobstructed singularity X_0 . Then M is modular if and only if $\tilde{T}^1(X/S) \otimes_{\mathcal{O}_S} \mathcal{O}_M$ is a flat \mathcal{O}_M -module.

Proof. There is the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & T^0(X/S) & \longrightarrow & \mathcal{O}_X^n & \xrightarrow{J(F)} & \mathcal{O}_X^k \longrightarrow \tilde{T}^1(X/S) \longrightarrow 0 \\ & & \downarrow \text{ev} & & \downarrow & & \downarrow \\ 0 & \longrightarrow & T^0(X_0) & \longrightarrow & \mathcal{O}_{X_0}^n & \xrightarrow{J(f)} & \mathcal{O}_{X_0}^k \longrightarrow T^1(X_0) \longrightarrow 0 \end{array}$$

Since X_0 was assumed to be unobstructed, \mathcal{O}_X is smooth and it follows that it is a free \mathcal{O}_S -module. Hence we can interpret $T^0(X/S)$ (resp. $T^0(X_0)$) as the syzygy module of the columns of the presentation matrix of $\tilde{T}^1(X/S)$ (resp. $T^1(X_0)$). Thus, by the lifting criterion for flatness (see [Mat86, § 7], for example), $\tilde{T}^1(X/S) \otimes_{\mathcal{O}_S} \mathcal{O}_M = \tilde{T}^1(X|_M/M)$ is a flat \mathcal{O}_M -module if and only if $ev|_M$ is surjective. \blacksquare

If X_0 is a complete intersection then $\tilde{T}^1(X/S)$ equals $T^1(X/S)$, hence:

Corollary 4.2. The modular stratum of an isolated complete intersection singularity X_0 equals the flattening stratum of the relative Tjurina module $T^1(X/S)$, where $\xi : X \rightarrow S$ is a semi-universal deformation of X_0 .

Remark. This result is extended to reduced space curve singularities in [Mar] using their determinantal structure (defining equations of any deformation of such a singularity are obtained as maximal minors of some $q \times (q-1)$ -matrix A , and the relative normal module of the deformation has a presentation matrix whose entries are the $(q-2)$ -minors of A). However it is not clear whether these assertions can be carried over to arbitrary isolated singularities, or at least to arbitrary unobstructed ones.

Nevertheless, analogous statements hold for deformations with section, which we are now going to derive:

Proposition 4.3. Let $\xi : X \rightarrow S$ be a semi-universal deformation of an unobstructed singularity X_0 with section $\sigma : S \rightarrow X$. Let X_0 be defined by $f_1, \dots, f_k \in \mathbb{C}\{x_1, \dots, x_n\}$, X be given by $F_1, \dots, F_k \in \mathcal{O}_S\{x_1, \dots, x_n\}$. Denote $J_\sigma := \text{Ker}(\sigma^*) \subseteq \mathcal{O}_X$, and let $J(F) := \left(\frac{\partial F_i}{\partial x_j} \right)_{i,j} \in \text{Mat}(k, n; \mathcal{O}_X)$ be the relative Jacobian matrix of F_1, \dots, F_k .

Then a subgerm $M \subseteq S$ is modular if and only if $\tilde{T}^1(X/S, J_\sigma) \otimes_{\mathcal{O}_S} \mathcal{O}_M$ is a flat \mathcal{O}_M -module, where

$$\tilde{T}^1(X/S, J_\sigma) := \mathcal{O}_X^k / J_\sigma \cdot J(F).$$

Proof. We may assume that $J_\sigma = (x)\mathcal{O}_X$. There is the following commutative diagram whose rows are exact sequences:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & T^0(X/S, J_\sigma) & \longrightarrow & \mathcal{O}_X^{n^2} & \xrightarrow{(x_1 J(F), \dots, x_n J(F))} & \mathcal{O}_X^k & \longrightarrow & \tilde{T}^1(X/S, J_\sigma) & \longrightarrow & 0 \\ & & \downarrow ev & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & T^0(X_0, \mathfrak{m}) & \longrightarrow & \mathcal{O}_{X_0}^{n^2} & \xrightarrow{(x_1 J(f), \dots, x_n J(f))} & \mathcal{O}_{X_0}^k & \longrightarrow & \mathcal{O}_{X_0}/(\mathfrak{m}J(f)) & \longrightarrow & 0. \end{array}$$

As in the proof of Proposition 4.1 we can interpret $T^0(X/S, J_\sigma)$ (resp. $T^0(X, \mathfrak{m})$) as syzygy modules of the modules $\tilde{T}^1(X/S, J_\sigma)$ (resp. $\mathcal{O}_{X_0}/(\mathfrak{m}J(f))$). So the surjectivity of $ev|_M$ means precisely that every syzygy of $\tilde{T}^1(X/S, J_\sigma)$ over the special fibre $s = 0$ lifts to a syzygy over M , i. e. $\tilde{T}^1(X/S, J_\sigma) \otimes_{\mathcal{O}_S} \mathcal{O}_M = \tilde{T}^1(X|_M/M, J_{\sigma|_M})$ is \mathcal{O}_M -flat. \blacksquare

Corollary 4.4. If X_0 is an isolated complete intersection, then the modular stratum (with section) of X_0 coincides with the flattening stratum of the relative Tjurina module $T^1(X/S, J_\sigma)$, where S is the base space of a semi-universal deformation $X \rightarrow S$ of X_0 and $J_\sigma = \text{ker}(\sigma^*) \subseteq \mathcal{O}_X$ corresponds to the section $\sigma : S \rightarrow X$.

Proof. X_0 being a complete intersection we have, for any subspace $M \subseteq S$, an exact sequence

$$0 \rightarrow T^1(X/S, J_\sigma)|_M \rightarrow \tilde{T}^1(X/S, J_\sigma)|_M \xrightarrow{r} \mathcal{O}_M \rightarrow 0$$

of \mathcal{O}_M -modules, where r is the residue map. Hence $T^1(X/S, J_\sigma)|_M$ is \mathcal{O}_M -flat if and only if $\tilde{T}^1(X/S, J_\sigma)|_M$ is. \blacksquare

These characterizations of modularity as flatness of a suitable \mathcal{O}_S -module make it possible to compute the modular stratum of X_0 , at least up to a given order. Computing the constant rank stratum or the annihilator of the associated torsion module whose vanishing implies flatness (see e.g. [GP02, section 7.3]) is not a feasible approach in this situation: The objects involved are too complicated for computations. Instead, one can determine, starting with the base field \mathbb{C} , the maximal small extension in \mathcal{O}_S that preserves flatness. The resulting algorithm and details on its implementation in the SINGULAR-library `modular.lib` ([Mar02c]) are described in [Mar02a], where also several non-trivial examples of modular strata for the functor Def_{X_0} are presented.

Remark. Of course one may consider the flattening stratum of $T^1(X/S)$ for any deformation $\xi : X \rightarrow S$ of X_0 . But if ξ is not a monodeformation, there is no connection to modular subspaces: The latter must be trivial, whereas the former may not, as the following example shows. Take an arbitrary hypersurface singularity X_0 defined by $f \in \mathcal{O}_n$ and $g \in (f, J(f))$. Then $f + \varepsilon g$ gives a trivial deformation X_D of X_0 over the double point D , its Kodaira-Spencer map is the zero map. On the other hand $T^1(X_D/D)$ is a flat $\mathcal{O}_D = \mathbb{C}[\varepsilon]$ -module:

Indeed, let $rf + \sum_k r_k \frac{\partial f}{\partial x_k} = 0$ be a relation of $f, J(f)$. Then $r(f + \varepsilon g) + \sum_k r_k \frac{\partial}{\partial x_k}(f + \varepsilon g) = \varepsilon(rg + \sum_k r_k \frac{\partial g}{\partial x_k})$, and one checks that this again belongs to $\varepsilon(f, J(f))$, so is equal to $\varepsilon(pf + \sum_k p_k \frac{\partial f}{\partial x_k})$ for some $p, p_k \in \mathcal{O}_n$. Then $(r - \varepsilon p, r_1 - \varepsilon p_1, \dots, r_n - \varepsilon p_n)$ lifts the given relation to $\mathcal{O}_n[\varepsilon]$. Hence any trivial first-order infinitesimal deformation of a hypersurface has flat relative $T^1(X_D/D)$.

4.2 Comparing flatness conditions

As established above, flatness of the module $T^1(X/S)$ (resp. $T^1(X/S, J_\sigma)$) has an interpretation as modularity of the corresponding deformation $\xi : X \rightarrow S$ (with section σ). One can then ask about the relationship to flatness of the 0-th cotangent cohomology module. The result is:

Proposition 4.5. Let $\xi : X \rightarrow S$ be a deformation of an isolated complete intersection singularity X_0 . If $T^1(X/S)|_M$ is a flat \mathcal{O}_M -module, then so is $T^0(X/S)|_M$, i. e. the flattening stratum of $T^1(X/S)$ is contained in the flattening stratum of $T^0(X/S)$, and this inclusion may be strict.

The analogous statement holds for deformations with section.

Proof. By splitting the short exact sequence

$$0 \rightarrow T^0(X/S)|_M \rightarrow \mathcal{O}_{X|_M}^n \rightarrow \mathcal{O}_{X|_M}^k \rightarrow T^1(X/S)|_M \simeq \tilde{T}^1(X|_M/M) \rightarrow 0$$

from the proof of Proposition 4.1 into two short exact sequences we deduce that $T^0(X/S)|_M$ is flat if $T^1(X/S)|_M$ is. In the case of a deformation with section we use the sequence in the proof of Proposition 4.3 instead.

On the other hand, let X_0 be a quasihomogeneous complete intersection defined by $f = (f_1, \dots, f_k) \in \mathcal{O}_n$ and take a basis monomial of $T^1(X_0)$ resp. $T^1(X_0, \mathfrak{m})$ which is of different weighted degree than f . Let $\xi : X_D \rightarrow D$ be the deformation over the double point D defined by $f + \varepsilon g$ (with the obvious section σ). From Proposition 3.8 we conclude that this monodeformation has trivial modular stratum, hence the flattening stratum of $T^1(X_D/D)$ resp. $T^1(X_D/D, J_\sigma)$ is trivial. But, as a submodule of $\mathcal{O}_{X_D}^n$ (resp. $\mathcal{O}_{X_D}^{n^2}$), $T^0(X_D/D)$ (resp. $T^0(X_D/D, J_\sigma)$) is a torsion-free, hence a flat $\mathcal{O}_D = \mathbb{C}[\varepsilon]$ -module. ■

We finish by comparing the modular strata of a hypersurface with respect to both deformation functors under consideration. First of all note the following fact:

Lemma 4.6. Let $\xi : X \rightarrow S$ be a deformation of X_0 with section $\sigma : S \rightarrow X$. If $M \subseteq S$ is modular with respect to the functor Def_{X_0} (i. e. considering ξ as an 'ordinary' deformation), then it is modular with respect to $Def_{X_0}^s$, too.

Proof. Suppose M is Def_{X_0} -modular. Let $\varphi : T \rightarrow M$ and $\psi : T \rightarrow S$ be morphisms such that $\varphi^*(\xi|_M)$ and $\psi^*(\xi)$ are isomorphic deformations with section of X_0 . Then they also give the same element of $Def_{X_0}(S)$, hence $\varphi = \psi$ by assumption. Thus M is $Def_{X_0}^s$ -modular, too. ■

We can prove a converse of this statement if we restrict ourselves to deformations with *singular section* of X_0 : If X_0 is equidimensional of dimension d , then a deformation with singular section of X_0 is a deformation $\xi : X \rightarrow S$ with section $\sigma : S \rightarrow X$, such that there is a factorization

$$\begin{array}{ccc} S & \xrightarrow{\sigma} & X, \\ & \searrow & \nearrow \\ & V(F^d(\Omega_{X/S})) & \end{array}$$

where $F^d(\Omega_{X/S})$ is the d -th Fitting ideal of the module of Kähler differentials of the \mathcal{O}_S -module \mathcal{O}_X , $C_\xi := V(F^d(\Omega_{X/S}))$ is the *critical locus* of ξ . One then has the following properties:

- (1) This definition gives rise to a subfunctor $Def_{X_0}^{ss}$ of $Def_{X_0}^s$. A versal deformation of X_0 with respect to $Def_{X_0}^{ss}$ is obtained as follows: Let $X \rightarrow S$ be a versal deformation of X_0 (with respect to Def_{X_0}), then, as seen in section 2, $X \times_S X \rightarrow X$ is versal for $Def_{X_0}^s$, and then the deformation induced by the base change $V(F^d(\Omega_{X/S})) \hookrightarrow X$ is a versal deformation with singular section of X_0 (cf. [Buc81, 2.5.1]).
- (2) If $f \in \mathcal{O}_n$ defines an isolated hypersurface singularity, then one can easily define a semi-universal deformation with singular section of $X_0 := (V(f), 0)$ by $F := f + \sum_i s_i g_i$, where $\{g_i\}_i$ represent a \mathbb{C} -basis of $\mathfrak{m} \cdot T^1(X_0, \mathfrak{m})$, \mathfrak{m} the maximal ideal of \mathcal{O}_{X_0} .

The latter construction cannot be generalized to singularities of higher codimension. For hypersurfaces, we can interpret deformations with singular section as deformations leaving the embedding dimension constant, the terms of its defining equation then belong \mathfrak{m}^2 . This is why we obtain the following result only for hypersurfaces.

Proposition 4.7. Let $\xi : X \rightarrow S$ be a deformation with singular section of a hypersurface X_0 and let $M \subseteq S$. If $T^1(X/S)|_M$ is a flat \mathcal{O}_M -module, then so is $T^1(X/S, J_\sigma)|_M$.

Proof. The short exact sequence of $\mathcal{O}_{X|M}$ -modules

$$0 \rightarrow J_{\sigma|M} \rightarrow \mathcal{O}_{X|M} \rightarrow \mathcal{O}_{X|M}/J_{\sigma|M} \simeq \mathcal{O}_M \rightarrow 0$$

yields the long exact cohomology sequence

$$\begin{aligned} 0 \longrightarrow T^0(X/S, J_{\sigma})|_M \xrightarrow{\iota_0} T^0(X/S)|_M \xrightarrow{\pi_0} T^0(X/S, \mathcal{O}_S)|_M \xrightarrow{\delta_0} \\ \xrightarrow{\delta_0} T^1(X/S, J_{\sigma})|_M \xrightarrow{\iota_1} T^1(X/S)|_M \xrightarrow{\pi_1} T^1(X/S, \mathcal{O}_S)|_M \longrightarrow 0. \end{aligned} \quad (4.1)$$

Let $f \in \mathcal{O}_n$ define X_0 . Since $X|M \rightarrow M$ is a deformation with singular section of X_0 , we may assume that $X|M$ is defined by $F \in (x)^2\mathcal{O}_M\{x\}$. In this situation we have $T^1(X/S, \mathcal{O}_S)|_M \simeq \mathcal{O}_M$ since it is the cokernel of

$$\begin{aligned} \mathrm{Der}_{\mathcal{O}_M}(\mathcal{O}_M\{x\}, \mathcal{O}_M) &\rightarrow \mathrm{Hom}_{\mathcal{O}_{X|M}}((F)/(F^2), \mathcal{O}_M) \simeq \mathcal{O}_M, \\ \delta &\mapsto \delta(F) \end{aligned}$$

which is the zero map as $F \in (x)^2\mathcal{O}_M\{x\}$. In addition, $\mathrm{Ker}(\iota_1)$ is the \mathcal{O}_M -module generated by $\frac{\partial F}{\partial x_1}, \dots, \frac{\partial F}{\partial x_n}$ modulo $(F) + (x_j \frac{\partial F}{\partial x_i})_{i,j}$. But this is a free \mathcal{O}_M -module: namely, suppose $\sum_i G_i(s) \frac{\partial F}{\partial x_i} = H \cdot F + \sum_{i,j} H_{ij} x_j \frac{\partial F}{\partial x_i}$. Since $(F, \frac{\partial F}{\partial x_1}, \dots, \frac{\partial F}{\partial x_n})$ form a regular sequence, this implies that, for all i , $G_i(s) - \sum_j H_{ij} x_j \in (F, \frac{\partial F}{\partial x_1}, \dots, \frac{\partial F}{\partial x_n})$, i. e.

$$G_i(s) = \sum_j H_{ij} x_j + A \cdot F + \sum_k B_k \frac{\partial F}{\partial x_k}$$

for some $A, B_k \in \mathcal{O}_M\{x\}$. But this is only possible if $G_i = 0$ since $F, \frac{\partial F}{\partial x_1}, \dots, \frac{\partial F}{\partial x_n} \in (x)\mathcal{O}_M\{x\}$.

Thus the long exact sequence (4.1) can be split into two short exact sequences

$$0 \rightarrow \mathrm{Ker}(\iota_1) \simeq \mathcal{O}_M \rightarrow T^1(X/S, J_{\sigma})|_M \rightarrow \mathrm{Im}(\iota_1) \rightarrow 0$$

and

$$0 \rightarrow \mathrm{Ker}(\pi_1) \rightarrow T^1(X/S)|_M \rightarrow \mathcal{O}_M \rightarrow 0,$$

from which we derive that if $T^1(X/S)|_M$ is \mathcal{O}_M -flat, then so is $T^1(X/S, J_{\sigma})|_M$. ■

Combining Lemma 4.6 and Proposition 4.7 with the modularity criteria Corollary 4.2 and Corollary 4.4, we obtain:

Corollary 4.8. Let $\xi : X \rightarrow S$ be a deformation with singular section $\sigma : S \rightarrow X$ of a hypersurface X_0 . Then

- (1) The flattening strata of $T^1(X/S)$ and $T^1(X/S, J_{\sigma})$ coincide.
- (2) If ξ is a monodeformation for both deformation functors Def_{X_0} and $\mathrm{Def}_{X_0}^s$, then the maximal modular subspaces inside S with respect to these two functors coincide.

References

- [Ale85] A. G. Aleksandrov, *Cohomology of a quasihomogeneous complete intersection*, Izv. Akad. Nauk SSSR Ser. Mat. **49** (1985), no. 3, 467–510, 672, *english translation*: Math. USSR-Izv. **49** (1985), no. 3, 437–477.
- [And74] M. André, *Homologie des algèbres commutatives*, Springer-Verlag, Berlin, 1974, Die Grundlehren der mathematischen Wissenschaften, Band 206.
- [Buc81] R. O. Buchweitz, *Contributions à la théorie des singularités*, Ph.D. thesis, Université Paris VII, 1981.
- [Eis95] D. Eisenbud, *Commutative Algebra with a View Toward Algebraic Geometry*, Graduate Texts in Mathematics, vol. 150, Springer-Verlag, New York, 1995.
- [Fle78] H. Flenner, *Über Deformationen holomorpher Abbildungen*, Habilitationsschrift, Universität Osnabrück, 1978.
- [GP02] G.-M. Greuel and G. Pfister, *A SINGULAR Introduction to Commutative Algebra*, Springer, Berlin, 2002.
- [GPS01] G.-M. Greuel, G. Pfister, and H. Schönemann, SINGULAR 2.0, A Computer Algebra System for Polynomial Computations, Centre for Computer Algebra, University of Kaiserslautern, 2001, <http://www.singular.uni-kl.de>.
- [Gra72] H. Grauert, *Über die Deformation isolierter Singularitäten analytischer Mengen*, Invent. Math. **15** (1972), 171–198.
- [KS90] S. Kosarew and H. Stieber, *A construction of maximal modular subspaces in local deformation theory*, Abh. Math. Sem. Univ. Hamburg **60** (1990), 17–36.
- [Lau79] O. A. Laudal, *Formal moduli of algebraic structures*, Lecture Notes in Mathematics, vol. 754, Springer, Berlin, 1979.
- [Loo84] E. J. N. Looijenga, *Isolated singular points on complete intersections*, London Mathematical Society Lecture Note Series, vol. 77, Cambridge University Press, Cambridge, 1984.
- [LP88] O. A. Laudal and G. Pfister, *Local moduli and singularities*, Lecture Notes in Mathematics, vol. 1310, Springer-Verlag, Berlin, 1988.
- [Mar] B. Martin, *Modular Deformations and Space Curve Singularities*, in: Algebraic Geometry and Singularities. Sevilla 2001, to appear in Rev. Mat. Iber.
- [Mar99] B. Martin, *Computing versal deformations with SINGULAR*, Algorithmic algebra and number theory (Heidelberg, 1997), Springer, Berlin, 1999, pp. 283–293.
- [Mar02a] B. Martin, *Algorithmic computation of flattenings and of modular deformations*, J. Symbolic Comput. **34** (2002), no. 3, 199–212.
- [Mar02b] B. Martin, *deform.lib*, a SINGULAR 2.0 library for computing miniversal deformations of singularities and modules, 2002.

- [Mar02c] B. Martin, `modular.lib`, a SINGULAR 2.0 library for computing the modular stratum, 2002.
- [Mat86] H. Matsumura, *Commutative ring theory*, Cambridge University Press, Cambridge, 1986.
- [MvS01] D. Mond and D. van Straten, *The structure of the discriminant of some space-curve singularities*, Q. J. Math. **52** (2001), no. 3, 355–365.
- [Pal76] V. P. Palamodov, *Deformations of Complex spaces*, Russian Mathematical Surveys **31** (1976), no. 3, 129–197.
- [Pal78] V. P. Palamodov, *Moduli in versal deformations of complex spaces*, Variétés analytiques compactes (Colloq., Nice, 1977), LNM 683, Springer, Berlin, 1978, pp. 74–115.
- [Pal82] V. P. Palamodov, *Cohomology of analytic algebras*, Trudy Moskov. Mat. Obshch. **44** (1982), 3–61, engl.: Trans. Moscow Math. Soc. 1983, no. 2 (44), 1–61.
- [Pal90] V. P. Palamodov, *Tangent fields on deformations of complex spaces*, Mat. Sb. **181** (1990), no. 10, 1320–1340.
- [Qui70] D. Quillen, *On the (co-) homology of commutative rings*, Applications of Categorical Algebra (Proc. Sympos. Pure Math., Vol. XVII, New York, 1968), Amer. Math. Soc., Providence, R.I., 1970, pp. 65–87.
- [Sai80] K. Saito, *Theory of logarithmic differential forms and logarithmic vector fields*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **27** (1980), no. 2, 265–291.
- [vS95] D. van Straten, *A note on the discriminant of a space curve*, Manuscripta Math. **87** (1995), no. 2, 167–177.